

CONTENTS

X-Ray investigation of benzyl 3,4-anhydro-2- <i>O</i> -mesyl-6- <i>O</i> -trityl- α -D-altropyranoside: ring conformations in the 3,4-anhydropyranose system J. W. Krajewski, P. Gluziński, and A. Banaszek (Warsaw, Poland)	1
Structural determination of alginic acid and the effects of calcium binding as determined by high-field n.m.r. C. A. Steginsky, J. M. Beale, H. G. Floss, and R. M. Mayer (Columbus, OH, U.S.A.) . .	11
PCIO quantum-mechanical relaxed conformational energy map of methyl 4-thio- α -maltoside in solution K. Mazeau (Grenoble, France) and I. Tvaroska (Bratislava, Czechoslovakia)	27
¹ H-N.m.r. spectral assignments for two series of heparin-derived oligosaccharides A. Horne and P. Gettins (Nashville, TN, U.S.A.)	43
Mode of formation of quinoxaline <i>versus</i> 2[1 <i>H</i>]-quinoxalinone rings from dehydro-D-erythorbic acid A. Mousaad, N. Rashed, H. A. Hamid, Y. El Kilany, and E. S. H. El Ashry (Alexandria, Egypt)	59
Spacer-modified saccharides for the regioselective photoaffinity labelling of the binding site of an immunoglobulin J. Lehmann and M. Scheuring (Freiburg, Germany)	67
Spacer-modified, photolabile tetrasaccharides as analogues of maltopentaose are versatile probes for porcine pancreatic α -amylase J. Lehmann and L. Ziser (Freiburg, Germany)	83
Synthesis, reactions, and crystal structure of 5-azido-6- <i>O</i> -benzoyl-3,5-dideoxy-3-fluoro-1,2- <i>O</i> -isopropylidene- α -D-glucopyranose C.-K. Lee, H. Jiang, and L. L. Koh (Kent Ridge, Singapore)	99
Esters boriques d'alditols. Synthèse, structure et stabilité en solution dans l'eau A. Munoz and L. Lamandé (Toulouse, France)	113
A convenient method for the preparation of a variety of ¹³ C-substituted D-fructose phosphates using readily available enzymes of the glycolytic pathway K. T. Kuan, D. S. Weber, L. Sottile, and W. J. Goux (Richardson, TX, U.S.A.)	123
<i>Notes</i>	
Analysis of glycosaminoglycan-derived oligosaccharides using fast-atom-bombardment mass-spectrometry R. J. Linhardt, H. M. Wang, D. Loganathan, D. J. Lamb, and L. M. Mallis (Iowa City, IA, U.S.A.)	137
Synthesis of a glycosidic affinity ligand for purification of cytidine-5'-monophosphosialate synthase J. Thiem, P. Stangier, P. Matschulat, and H. Paulsen (Hamburg, Germany)	147
Synthesis of 1,6-anhydro-2,3-dideoxy- β -D-erythro- and -threo-hex-2-enopyranose from 3,4-di- <i>O</i> -substituted glycals by a facile intramolecular Ferrier reaction H. B. Meryyala, K. C. Venkataramanaiah, and V. S. Dalvoy (Hyderabad, India)	151

A synthesis of 2,3-dideoxy-4,6:7,8-di- <i>O</i> -isopropylidene-D- <i>gluco</i> -oct-2-enono-1,5-lactone S. Valverde, S. Garcia-Ochoa, and A. M. Gomez (Madrid, Spain)	155
Reaction of Meldrum's acid with D-mannose and L-arabinose F. Z. Mata, M. B. Martínez, and J. A. Galbis Pérez (Sevilla, Spain)	159
Synthesis of <i>p</i> -trifluoroacetamidophenyl 2-acetamido-4- <i>O</i> -(2-acetamido-2-deoxy- β -D-glucopyranosyl)-2-deoxy- β -D-mannopyranosiduronic acid, an artificial antigen corresponding to a disaccharide repeating unit of the capsular polysaccharide of <i>Haemophilus influenzae</i> type e P. J. Garegg, S. Oscarson, and A.-K. Tidén (Stockholm, Sweden)	163
Structure of the acidic exopolysaccharide secreted by <i>Rhizobium leguminosarum</i> biovar. <i>phaseoli</i> CFN42 A. Gil-Serrano, I. González-Jiménez, P. Tejero-Mateo, A. Sánchez del Junco, M. Megias, and M. J. Romero-Vázquez (Sevilla, Spain)	169
Some <i>N</i> -acyl derivatives of <i>O</i> -carboxymethylchitosan S. Hirano, K. Hayashi, and K. Hirochi (Tottori, Japan)	175
<i>Papers</i>	
Synthesis of aryl glycosides as <i>vir</i> gene inducers of <i>Agrobacterium tumefaciens</i> D. Delay, J. Cizeau, and F. Delmotte (Orléans, France)	179
Synthesis of 3,6-dideoxy-4- <i>C</i> -(4 ¹ -hydroxyethyl)hexopyranoses (yersinioses) from 1,6-anhydro- β -D-glucopyranose V. A. Zubkov, R. P. Gorshkova, Yu. S. Ovodov (Vladivostok, U.S.S.R.), A. F. Sviridov, and A. S. Shashkov (Moscow, U.S.S.R.)	189
The preparation, resolution, and phosphorylation of some benzyl ethers of <i>myo</i> -inositol: intermediates for the synthesis of <i>myo</i> -inositol phosphates of the phosphatidylinositol cycle T. Desai, J. Gigg, and S. Payne (London, Gt. Britain)	209
Synthesis of trisaccharide methyl glycosides related to fragments of the capsular polysaccharide of <i>Streptococcus pneumoniae</i> type 18C A. M. P. van Steijn, J. P. Kamerling, and J. F. G. Vliegenthart (Utrecht, The Netherlands)	229
Synthesis of 2,6,7-trideoxy-7- <i>C</i> -(2,4-dichlorophenyl)-D- <i>xylo</i> -heptonic acid and 6-(2,4-dichlorophenyl)-D- <i>xylo</i> -2,3,4-trihydroxyhexanesulfonic acid J. Kuszmán and B. Podányi (Budapest, Hungary)	247
Synthesis of 7-(2,4-dichlorophenyl)-D- <i>erythro</i> -3-hydroxy-5-heptanolide, 6-(2,4-dichlorophenyl)-D- <i>erythro</i> -2,4-dihydroxyhexane-1-sulfonic acid, and 6-(2,4-dichlorophenyl)-D- <i>erythro</i> -2,4-dihydroxyhexylphosphonic acid G. Hodosi, G. Galambos, B. Podányi, and J. Kuszmán (Budapest, Hungary)	269
Synthesis of lysine-containing fragments of the <i>Proteus mirabilis</i> O27 O-specific polysaccharide and neoglycoconjugates therefrom A. Ya. Chernyak, L. O. Kononov, P. Radha Krishna, N. K. Kochetkov (Moscow, U.S.S.R.), and A. V. Rama Rao (Hyderabad, India)	279
Preparation and characterization of <i>N</i> -[2-(glycosyloxy)ethyl]chitosan derivatives K. R. Holme and L. D. Hall (Vancouver, B.C., Canada)	291
Solution interaction and rheology of <i>N</i> -[2-(glycosyloxy)ethyl]chitosan derivatives K. R. Holme, L. D. Hall, R. A. Speers, and M. A. Tung (Vancouver, B.C., Canada)	307

Inhibition of dextran synthesis by acceptor reactions of dextransucrase, and the demonstration of a separate acceptor binding-site A. Tanriseven and J. F. Robyt (Ames, IA, U.S.A.)	321
<i>Note</i>	
Glucuronoxylomannan of <i>Cryptococcus neoformans</i> serotype C: structural analysis by gas-liquid chromatography-mass spectrometry and ^{13}C -nuclear magnetic resonance spectroscopy R. Cherniak, L. C. Morris, and S. A. Meyer (Atlanta, GA, U.S.A.)	331
<i>Author index</i>	c1
<i>Subject index</i>	c3
<i>Contents (vol. 225)</i>	c7